

[Skip to main content](#) [Skip to sections](#)

This service is more advanced with JavaScript available, learn more at <http://activatejavascript.org>

[Search](#)

- [Home](#)
- [Contact Us](#)
- [Log in](#)

Search

[Download PDF](#)



[Journal of the Korean Society for Applied Biological Chemistry](#)

December 2015, Volume 58, [Issue 6](#), pp 839–846

Bioactive carbazole alkaloids from *Alysicarpus ovalifolius* (Schumach)

- Authors
- [Authors and affiliations](#)

- Angelina Atieno Ochung' [Email author](#)
- Lawrence Arot Oyango Manguro
- Phillip Okinda Owuor
- Isaac Ogoche Jondiko
- Regina Achieng' Nyunja
- Hosea Akala
- Pauline Mwinzi

- Sylvia Awino Opiyo

Article

First Online:

19 August 2015

Received:

15 May 2015

Accepted:

24 June 2015

DOI: 10.1007/s13765-015-0100-4

Cite this article as:

Ochung', A.A., Manguro, L.A.O., Owuor, P.O. et al. J Korean Soc Appl Biol Chem (2015) 58: 839. doi:10.1007/s13765-015-0100-4

- 64 Downloads

Abstract

Phytochemical and biological evaluation of the stem bark of *Alysicarpus ovalifolius* led to the isolation of three carbazole alkaloids identified as mohanimbine (**1**), koenimbine (**2**) and koenidine (**3**) along with quercetin 3-*O*-glucoside (**4**), kaempferol 7-*O*-glucoside (**5**), orientin (**6**), apigenin (**7**), quercetin (**8**), plumbagin (**9**) and stigmasterol (**10**). The structures of these compounds were elucidated using physical and spectroscopic methods as well as comparison with the literature data. Compound **3** showed strong activity against chloroquine-sensitive strain I (D6) and the multi-drug resistant Indochicha I (W2) of *Plasmodium falciparum* with IC₅₀ values of 63.07 ± 0.01 and 54.19 ± 0.04 ng/mL, respectively. Compound **1** on the other hand exhibited moderate larvicidal against *Anopheles gambiae* larvae as well as antimicrobial activities against *Candida albicans* and gram-positive *Staphylococcus aureus*, respectively.

Keywords

Alkaloids *Alysicarpus ovalifolius* Antiplasmodial Larvicidal Mosquitocidal Fabaceae

Introduction

The family Fabaceae consists of approximately 20,000 species spread in 650 genera and is rich in flavonoids, anthraquinones, alkaloids, terpenoids, lipids, and polysaccharides some of which have medicinal properties (Wojciechowski et al. 2004; Yenesew et al. 2004). The family is one of the most economically important plants in the provision of medicines, ornaments, dyes, timber, fodder, tannins, resins, essential oils, flavours, insecticides, piscicides, and even human food (Bentjee 1994; Mannetje 2002). The hitherto phytochemically uninvestigated *Alysicarpus ovalifolius* is used in folklore medicine in management of fever, wounds, ringworm, acute and chronic-troubled bleeding piles, and as a stimulant in birth control while leaf sap and root

decoction are drunk to relieve cough (Lamers et al. [1996](#); Kokwaro [2009](#)). In the current communication, phytochemical analysis of *n*-hexane, CH₂Cl₂ and MeOH extracts resulted in the isolation of three carbazole alkaloids (**1–3**) together with compounds (**4–10**). These compounds together with their activities are being reported from this plant for the first time.

Materials and methods

Experimentation, solvents and fine consumables

Melting points were determined using Gallenkamp melting point apparatus and are uncorrected. The NMR data were measured in CDCl₃ and CDCl₃-DMSO-d₆ on a JOEL NMR instrument operating 600 and 150 MHz, respectively. Some NMR analyses were done using Bruker AM 300 spectrometer operating at 300 and 75 MHz, respectively. TMS was used as internal standard. The mass spectral data were obtained using a Varian MAT 8200 A instrument. Column chromatography was performed using silica gel 60 (0.063–0.200 mm, Merck-Germany), while thin layer chromatography (TLC) was performed using silica gel 60 F₂₅₄ (Merck) pre-coated plates. All solvents used were of analytical grade.

The stem bark of *Alysicarpus ovalifolius* was obtained from Shimba Hills (latitude: 4°19'39"S and longitude: 39°21'39"E) where it naturally grows as a wild plant. The plant materials were authenticated at the herbarium of the Museums of Kenya where voucher specimen (No: FAB/AO/2012) was preserved.

Extraction and isolation

The air dried and pulverized root bark (1 kg) of the plant was soaked sequentially in *n*-hexane (3 × 3 L), CH₂Cl₂ (3 × 3 L) and MeOH (3 × 3 L), each lasting 4 days at room temperature. The extracts were separately filtered and evaporated under reduced pressure to afford yellowish (7 g), brown (25 g) and reddish-brown (176 g) extracts of *n*-hexane, CH₂Cl₂ and MeOH, respectively. Five gram of *n*-hexane extract was mixed with silica gel in a minimum amount of dichloromethane and chromatographed over silica gel-packed column (2.0 × 60 cm, SiO₂ 120 g) using *n*-hexane with increasing amount of CH₂Cl₂ up to 100 % of the latter. A total of 90 fractions, each 20 mL was collected and their homogeneity monitored by TLC (solvent systems: *n*-hexane EtOAc, 9:1 and 4:1). The eluants were grouped into three pools (I-III) depending on TLC profiles. Fractions 1–25 constituted pool I, which upon evaporation of solvent afforded a yellow oily compound that lost colour with time and was discarded. Pool II (fractions 30–55, 1.5 g) showed a single yellow spot of *R_f* 0.61 (solvent system: *n*-hexane–EtOAc, 4:1) which on further recrystallization gave **10** (90 mg). Pool III (fractions 57–85, 1.0 g) also crystallized out to give a colourless powder which on further recrystallization afforded **9** (75 mg). Dichloromethane extract (22 g) was adsorbed onto silica gel and then subjected to column chromatography (2.5 × 60 cm, SiO₂ 240 g, pressure ≈ 1 bar) using *n*-hexane–CH₂Cl₂ gradient (increment 10 %) up to 100 % CH₂Cl₂ and elution concluded with ethyl acetate, collecting 20 mL each. The process afforded various sub-fractions (I–V) as determined by TLC profiles [solvent systems: *n*-hexane–CH₂Cl₂ (1:2, 1:1) and CH₂Cl₂–MeOH, 98:2 and 96:4]. The sub-fraction I (fractions 1–15) showed no spot and solvent recovered. Sub-fraction II (fractions 20–60) produced a single spot *R_f* 0.43 (eluent: *n*-hexane–CH₂Cl₂, 1:1) that was Dragendorff's reagent positive which upon

evaporation of solvent followed by crystallization in CH₂Cl₂–MeOH mixture afforded **1** (45 mg). Sub-fraction III (fractions 63–86, 3.7 g) afforded compounds of *R_f* values 0.43 and 0.38 and upon repeated chromatographic separation afforded a further **1** (15 mg) and **2** (20 mg), respectively. Sub-fraction IV (fraction 93–130, 3.4 g) showed one major spot *R_f* 0.60 (solvent system: CH₂Cl₂–MeOH, 97:3) and upon evaporation of solvent, followed by crystallization gave compound **2** in 165 mg. Fractions 134–180 constituted sub-fraction V (5 g) and was further purified by medium pressure chromatography (2.5 × 50 cm, SiO₂ 150 g, pressure ≈1 bar) to give further compounds **2** and **3** in 50 and 70 mg, respectively.

Medium pressure chromatographic separation of MeOH extract (150 g) over 2 % oxalic acid solution-deactivated silica gel column using a mixture of CH₂Cl₂–methanol (5 % increment of MeOH) and MeOH neat gave a total of 80 fractions of 50 mL each. Fractions exhibiting similar TLC profiles were pooled together (Pools A–C). Pool A (fractions 7–20, 5 g) showed one major spot *R_f* 0.33 (eluant: CH₂Cl₂–MeOH, 97:3) along with minor ones and was further purified by crystallization to give **8** (180 mg). Fractions 25–45 (pool B, 7 g) was repeatedly fractionated over 2 % oxalic acid-deactivated silica gel (SiO₂ 150 g; 2.5 × 50 cm; 2–3 % MeOH–CH₂Cl₂) affording sub-fractions which resulted into **8** (15 mg), **7** (35 mg) and **6** (70 mg). On the other hand, fraction C (5 g) yielded **5** (54 mg) and **4** (35 mg) under similar purification procedure.

Compound 1, colourless powder, mp 92–94 °C [lit. 88–90 °C (Abu Bakar et al. 2007)]

¹H NMR (CDCl₃, 300 MHz), δ 7.69 (4-H, 1H, s), 7.96 (5-H, 1H, d, *J* = 7.5 Hz), 7.19 (6-H, 1H, d, *J* = 7.5 Hz), 7.28 (7-H, 1H, d, *J* = 5.0 Hz), 7.36 (8-H, 1H, d, *J* = 7.2 Hz), 6.64 (3'-H, 1H, d, *J* = 9.6 Hz), 5.67 (4'-H, 1H, d, *J* = 9.6 Hz), 1.77 (5'-H, 2H, t, *J* = 8.1 Hz), 1.83 (6'-H, 2H, m), 5.13 (7'-H, 1H, t, *J* = 6.6 Hz), 1.62 (8'-H, s), 1.77 (9'-H, s), 1.49 (2'-CH₃, 3H, s), 2.38 (3-CH₃, 3H, s), 7.41 (1H, N–H, s).

¹³C NMR (CDCl₃, 75 MHz), δ 104.2 (C-1), 149.9 (C-2), 131.7 (C-3), 117.5 (C-4), 119.5 (C-5), 119.3 (C-6), 121.2 (C-7), 110.4 (C-8), 134.9 (C-9), 139.5 (C-10), 118.4 (C-11), 116.7 (C-12), 78.2 (C-2'), 128.5 (C-3'), 124.2 (C-4'), 40.8 (C-5'), 22.7 (C-6'), 116.7 (C-7'), 131.7 (C-8'), 25.9 (C-9'), 25.7 (C-10'), 17.6 (2'-CH₃). IR *v*_{max} (KBr) cm⁻¹: 3350, 2930, 2850, 1650, 1450, 1380, 1210, 760; 28¹H and ¹³C NMR (CDCl₃) ppm: see Tables 1 and 2; ESI-MS (rel. int): *m/z* 332.3 [M]⁺ (100), 331.3 (10), 276.2 (7), 250 (30), 248 (10), 210 (10).

Table 1

In vitro antiplasmodial activity (IC₅₀) of crude extracts *A. ovalifolius* root bark and the pure isolates against D6 and W2 strains of *Plasmodium falciparum*

Test materials	IC ₅₀ values ^a (ng/mL)	
	W2 Clone	D6 Clone
<i>n</i> -Hexane extract	323.0 ± 0.01	546.0 ± 0.01
Dichloromethane extract	234.0 ± 0.01	482.0 ± 0.03
Methanol extract	265.0 ± 0.04	579.0 ± 0.02
Pure isolates		
Mohanimbine (1)	130.0 ± 0.01	279.0 ± 0.01

Test materials	IC ₅₀ values ^a (ng/mL)	
	W2 Clone	D6 Clone
Koenimbine (2)	161.4 ± 0.02	176.7 ± 0.01
Koenidine (3)	63.0 ± 0.01	54.2 ± 0.04
Chloroquine (20 mg/mL)	17.5 ± 0.01	26.9 ± 0.01

^aValues are mean ± SD of three replicates

Table 2

Larvicidal and mosquitocidal activities of crude extracts and pure isolates as % mortality and LC₅₀ values

Test materials	Larvicidal activity		Mosquitocidal activity	
	Mortality ^a (%)	LC ₅₀ (µg/mL)	Mortality ^a (%)	LC ₅₀ (µg/mL)
<i>n</i> -Hexane extract	25.9 ± 0.01	120.61	5.1 ± 0.01	500.56
Dichloromethane extract	87.7 ± 0.01	9.86	88.5 ± 0.01	17.83
Methanol	77.8 ± 0.01	85.64	5.1 ± 0.01	435.17
Pure isolates				
Mohanimbine (1)	82.3 ± 0.01	5.56	44.3 ± 0.01	213.90
Koenimbine (2)	53.6 ± 0.01	41.76	42.6 ± 0.01	260.46
Koenidine (3)	34.7 ± 0.01	87.54	30.7 ± 0.1	289.07

^a Values are mean ± SD of three replicates recorded at a concentration of 250 µg/mL

Compound 2, colourless powder, mp. 196–197 °C [lit. 194–195 °C (Nayak et al. [2010](#))]

¹H NMR (CDCl₃, 300 MHz), δ 7.61 (4-H, 1H, s), 7.41 (5-H, 1H, d, *J* = 7.6 Hz), 7.25 (6-H, 1H, d, *J* = 7.2 Hz), 7.42 (8-H, 1H, d, *J* = 7.6 Hz), 6.78 (3'-H, 1H, d, *J* = 11 Hz), 5.68 (4'-H, 1H, d, *J* = 14.7 Hz), 1.47 and 2.28 (2 × 2'-CH₃, 3H, s), 2.29 (3-CH₃, 3H, s), 3.87 (7-OCH₃, s), 7.61 (1H, N-H, s).

¹³C NMR (CDCl₃, 75 MHz), 103.7 (C-1), 150.9 (C-2), 125.3 (C-3), 122.1 (C-4), 103.7 (C-5), 114.2 (C-6), 155.2 (C-7), 112.3 (C-8), 136.6 (C-9), 132.9 (C-10), 118.5 (C-11), 116.0 (C-12), 77.1 (C-2'), 130.1 (C-3'), 118.1 (C-4'), 28.2, 16.5 (C-2'), 56.7 (6-OCH₃). IR ν_{max} (KBr) cm⁻¹: 3330, 2960, 2870, 1642, 1470, 1363, 1312, 1208, 690; ²⁸H and ¹³C NMR (CDCl₃) ppm: see Tables [1](#) and [2](#); ESI-MS (rel. int): *m/z* 294.3 [M]⁺ (100), 296 (10), 293.3 (24).

Compound 3, pale yellow crystals, mp 225–226 °C [lit. 224–225 °C (Mohammad et al. [2013](#))]

^1H NMR (CDCl_3 , 300 MHz), δ 7.40 (4-H, 1H, s), 7.53 (5-H, 1H, s), 6.95 (8-H, 1H, s), 6.64 (3'-H, 1H, d, $J = 9.6$ Hz), 5.69 (4'-H, 1H, d, $J = 9.6$ Hz), 1.28 and 1.50 ($2 \times 2'$ - CH_3 , 3H, s), 2.35 (3- CH_3 , 3H, s), 3.99 (6- OCH_3 , s), 3.97 (7- OCH_3 , s), 7.30 (1H, N-H, s).

^{13}C NMR (CDCl_3 , 75 MHz), 103.4 (C-1), 146.7 (C-2), 133.9 (C-3), 115.7 (C-4), 101.1 (C-5), 142.5 (C-6), 146.9 (C-7), 93.7 (C-8), 133.9 (C-9), 117.1 (C-10), 117.1 (C-12), 74.2, (C-2'), 127.4 (C-3'), 118.7 (C-4'), 14.8 and 26.3 ($2 \times 2'$ - CH_3), 55.3 (6- OCH_3), 54.8 (7- OCH_3). IR ν_{max} (KBr) cm^{-1} : 3150, 2940, 2880, 1640, 1450, 1344, 1110, 860; ^{28}H and ^{13}C NMR (CDCl_3) ppm: see Tables [1](#) and [2](#); ESI-MS (rel. int): m/z 324.3 [M] $^+$ (100), 323.3 (50), 308 (5), 282 (5).

Acid hydrolysis of compounds 4 and 5

A solution of **4** and **5** (each 10 mg) in a mixture of 8 % HCl (1 mL) and MeOH (20 mL) was separately refluxed for 2 h. The reaction mixtures were reduced in vacuo to dryness, dissolved in H_2O (2 mL) and neutralized with NaOH. The neutralized products were then subjected to TLC analysis (eluent: EtOAc–MeOH– H_2O –HOAc, 6:2:1:1). The chromatograms were sprayed with aniline hydrogen phthalate followed by heating at 100 °C for 2 min. The presence of glucose was confirmed after comparison with authentic samples.

Antiplasmodial assay

An in vitro antiplasmodial activity was carried out using the *P. falciparum* multi-drug resistant Indochicha I (W2) and chloroquine-sensitive Sierra Leone I (D6) strains according to procedures of Desjardins et al. ([1979](#)) and Chulay ([1983](#)) to determine the IC_{50} . The parasites were grown in a continuous culture supplemented with mixed gas (90 % nitrogen, 5 % oxygen), 10 % human serum and 6 % haematocrit of A+ red blood cell (Trager and Jensen [1976](#)). When the cultures had reached a parasitemia of 3 % with at least a 70 % ring developmental stage present, parasites were transferred to a 96-well microtitre plate with wells pre-coated with the extract/isolates. The samples were diluted across the plate to provide a range of concentrations used to accurately determine IC_{50} values. The plates were then incubated in a mixed gas incubator for 24 h, then 3H-hypoxanthine was added and the parasite allowed to grow for 18 h more in triplicates. Cells were then processed with a plate harvester (TomTec) on a filter paper and washed to eliminate unincorporated radioisotope. Chloroquine was used as a standard drug.

Larvicidal and mosquitocidal tests

Batches of 20 third instar *Anopheles gambiae* larvae were transferred by means of droppers to a small disposable test cup each containing 100 mL of five concentrations of the isolates of 0, 10, 100, 250, 500 and 1000 $\mu\text{g}/\text{mL}$ dissolved in acetone in triplicate vials according to (Globade et al. [2002](#)). Acetone was used as a positive control, while distilled water acted as a negative control. Dead larvae were counted after 24 h. For insecticidal bioassays, sample solutions of 0, 10, 100, 250, 500 and 1000 $\mu\text{g}/\text{mL}$ dissolved in acetone in triplicate vials were applied into several filter paper discs (5 cm diameter) then placed in perforated dishes (treated set). After drying the filter papers, ten unfed insects were introduced into each of the dishes through a hole and allowed to be in contact with the filter papers discs for 20 min, then transferred into cages with sugar water and observed for 24 h in triplicates (Gbolade et al. [2002](#)). The mortality of the


insects was monitored and toxicity levels of the test samples evaluated graphically to give LC50 values. Controls were distilled H₂O and solvent acetone.

Antifungal and antibacterial tests

The disc diffusion assay method was applied according to Singh, et al. [2002](#) using *Candida albicans* (HG 392), *Aspergillus fumigatus* (HG 420) and *Aspergillus niger* (ATCC 90028) as the representative fungi. *Staphylococcus aureus* (ATCC 25922), *Streptococcus faecalis* (ATCC 25925) and *Bacillus anthracis* (QST 713) were used as the representative gram-positive bacteria, while *Klebsiella pneumoniae* (ATCC 90028), *Salmonella typhimurium* (ATCC 25927), *Pseudomonas aeruginosa* (ATCC 25923) and *E. coli* (K 12) were representatives of gram-negative bacteria. Crude samples were tested in vitro at sample concentration of 1000 µg/mL dissolved in dimethylsulfoxide (DMSO) in triplicate vials, while pure isolates were tested at 100 µg/mL. Mueller–Hinton agar was aseptically aliquoted at volumes of 25 mL to Petri dishes and left to congeal. The agar was inoculated aseptically with test organisms using streaking method (Singh et al. [2002](#)). Test discs (5 mm diameter) previously impregnated with 10 µL of test samples were placed approximately equidistant into the seeded agar using a sterile forceps. Disc containing 10 µL of the 20 µg/mL solution of the standard drug (Fluconazole and Amoxicillin) was used as the positive control. The agar plates were incubated at 37 °C for 72 h after which the inhibition zones were measured in millimeters (McChesney et al. [1991](#)).

Results and discussion

Column chromatography separation of the CH₂Cl₂ extract (29 g) yielded three compounds (**1–3**), Fig. [1](#), which showed positive dragendorff's test for alkaloids. Compound **1** was isolated as colourless powder with a molecular formula C₂₃H₂₅O as evidenced by a molecular ion peak at m/z 332.3 [m]⁺. It showed a positive Dragendorff's test suggesting it is an alkaloid compound. The ¹H NMR spectrum of **1** (Fig. [2](#)) showed four methyl singlet signals which appeared at δ 1.77 and 1.62 (3H each, vinyl methyls), 2.38 (3H, an aromatic methyl) and 1.49 (3H) besides a vinyl proton on a trisubstituted double bond at δ 5.13 (br t, $J = 6.6$ Hz). The latter peak correlated with the vinyl methyls in the HMBC spectrum suggesting the presence of a terminal –CH₂CH=C(CH₃)₂ group in the compound (Abu Bakar et al. [2007](#); Mohammad et al. [2013](#)). In the ¹H NMR spectrum, a set of olefinic protons at C-3' and C-4' appeared as doublets at δ 6.64 ($J = 9.6$ Hz) and 5.67 ($J = 9.6$ Hz) which together with a ¹³C NMR (Fig. [3](#)) diagnostic peak at δ 78.2 signified the presence of carbon–carbon double bond in a pyran ring typical of carbazole alkaloids (Dheeref et al. [2014](#)). Comparison of ¹H and ¹³C NMR spectra of compound **1** with those of mohanimbine previously isolated from *Murraya koenigii* revealed close similarities (Chakraborty et al. [1978](#); Fiebig et al. [1985](#)). Analysis of both ¹H and ¹³C NMR data of compound **1** taking into consideration the fragmentation pattern in the ESI-MS confirmed the compound to be mohanimbine.



- 1 R = R' = H, R'' = CH₃, R''' = CH₂-CH₂-CH=C(CH₃)₃
- 2 R = H, R' = OCH₃, R'' = R''' = CH₃
- 3 R = R' = OCH₃, R'' = R''' = CH₃

Fig. 1

Structures of compounds **1**, **2** and **3**






Fig. 2

¹H NMR spectrum for compounds 1, 2 and 3






Fig. 3

¹³C NMR spectra for compounds **1**, **2** and **3**

Compound **2**, a colourless powder also afforded a positive Drangedorff's test for alkaloids. The ESI-MS molecular ion peak at m/z 294.3 and the ¹³C NMR data in combination with DEPT (90° and 135°) suggested a molecular formula C₁₉H₂₀NO₂ as deduced by comparative analysis of ¹³C NMR (19 distinct signals); DEPT-¹³C NMR (90°: 6CH all SP²); $\theta = 135^\circ$: 6CH and 4CH₃ including a methoxy group at δ 56.7. The DEPT data comprising of 6CH and 4CH₃ which totalled to C₁₀H₉ suggested the presence of either a hydroxyl or a –NH group in the compound. The ¹H NMR of compound **2** showed a set of olefinic signals at δ 6.78 (d, $J = 11.0$ Hz) and 5.68 (d, $J = 14.7$ Hz) which together with a broad singlet at δ 7.61 suggested that compound **2** is a carbazole alkaloid (Dheeref et al. [2014](#)). Comparative analysis of both ¹H and ¹³C NMR data of the compound taking into consideration of the molecular ion m/z 294.3, and the fragmentation pattern in the ESI-MS revealed the compound to be koenimbine previously isolated from *M. koenigii* (Ito et al. [2006](#); Mohammed et al. [2013](#)). On the basis of physical and spectroscopic data, compound **2** was confirmed to be koenimbine.

Compound **3** was obtained as pale yellow needles and the ESI-MS indicated a molecular ion peak at m/z 324.3 suggesting a molecular formula of C₂₀H₂₁NO₃ which is a 29 amu higher than that for compound **2**, thus suggesting that compound **3** is a derivative **2**. Comparison of the ¹H and ¹³C NMR spectra of **3** with those of **2** (Figs. [2](#) and [3](#)) showed close similarities with notable difference between the two compounds being the presence of additional methoxy group in **3** as evidenced by a ¹H NMR peak at δ 3.97 (s, 3H) with corresponding δ_C 54.8. Further confirmation of the structure of **3** was accomplished using HSQC and HMBC data which aided unequivocal assignment of all the signals associated with the compound. Thus, based on spectroscopic data coupled with reported literature data, compound **3** was established to be koenidine also previously isolated from *M. koenigii*.

The extracts and isolates (**1–3**) showed strong to moderate in vitro antiplasmodial activities against D6 and W2 strains as shown in Table [1](#). The lowest IC₅₀ values of 234.0 ± 0.01 and 63.0 ± 0.01 ng/mL were observed for the dichloromethane extract and koenidine (**3**), respectively, for the W2 strain of *P. falciparum* compared with the standard drug (chloroquine) whose value was 17.5 ± 0.01 ng/mL. Compound **3** also exhibited strong activity against the D6 strain of *P. falciparum* with IC₅₀ value of 54.2 ± 0.04 ng/mL compared with the standard drug (chloroquine) whose value was 26.9 ± 0.01 ng/mL. Thus, the strong activity of the dichloromethane extract could be attributed to the presence of the compound **3**. No antiplasmodial activity has been reported for the extracts/isolates.

The dichloromethane extract also exhibited the highest larvicidal activity at a concentration of 250 μ g/mL with 87.7 ± 0.01 % mortality recorded by the first time interval of 24 h. The methanol extract was equally active with mortality of 77.8 ± 0.01 %. Compound **1** was the most active of the pure isolated with 82.3 ± 0.01 % mortality. The lowest LC₅₀ values of 9.86 and 5.56 μ g/mL were observed for dichloromethane extract and compound **1**, respectively. Results are summarized in Table [2](#). The dichloromethane extract had the highest mosquitocidal activity of 88.5 ± 0.01 % mortality at a concentration of 250 μ g/mL after 100 min, while compound **1** showed 44.3 ± 0.01 % mortality with LC₅₀ values of 17.83 and 213.90 mg/mL, respectively:

The crude extracts and pure isolates were subjected to in vitro antifungal and antibacterial activities against the yeast-like and filamentous fungi as well as some gram-positive and gram-negative bacteria using the disc diffusion method. Results are presented in Table 3. Again the dichloromethane extract showed strongest activity against *C. albicans* and *S. aureus* with zones of inhibition measuring 13.2 ± 0.1 and 15.3 ± 0.1 mm, respectively, compared to Fluconazole and Amoxicillin, which were used as the standard drugs with zones of inhibition of 17.3 ± 0.2 and 19.5 ± 0.1 mm. Methanol extract was most toxic to *S. typhimurium* and *S. aureus* with zones of inhibition of 13.3 ± 0.1 and 12.3 ± 0.1 mm, respectively. Mohanimbine (**1**) exhibited strongest inhibition against *C. albicans* and *S. aureus* (14.5 ± 0.1 and 13.8 ± 0.1 mm, respectively). These results concur with the previous investigations (Mohammad et al. 2013, Dheeraj et al. 2014) which reported the antimicrobial activities of compounds **1**, **2** and **3** isolated from *M. koenigii*. Antimicrobial activities of some *Alysicarpus* species were previously reported (Rameshkumar and Umarajan 2013; Kumar et al. 2014). The results from this study confirm the ethnomedicinal information of plant.

Table 3

In vitro antifungal and antibacterial activities of extracts and pure isolates of *A. ovalifolius*

Test microbe	Diameter of zones of inhibition* in mm							
	N-hex	DCM	MeOH	Cpd 1	Cpd 2	Cpd 3	Flu	Am
<i>C. albicans</i>	2.3 ± 0.01	13.2 ± 0.1	9.1 ± 0.1	14.5 ± 0.1	8.0 ± 0.2	10.4 ± 0.2	17.3 ± 0.2	–
<i>A. fumigatus</i>	5.2 ± 0.01	7.4 ± 0.1	7.2 ± 0.2	6.7 ± 0.1	6.5 ± 0.2	7.3 ± 0.3	19.5 ± 0.1	–
<i>A. niger</i>	ND	5.8 ± 0.2	6.5 ± 0.1	6.5 ± 0.1	7.1 ± 0.3	8.5 ± 0.4	15.8 ± 0.3	–
<i>S. aureus</i>	7.2 ± 0.01	15.3 ± 0.1	12.3 ± 0.1	13.8 ± 0.1	6.4 ± 0.0	5.2 ± 0.1	–	19.5 ± 0.1
<i>S. faecalis</i>	4.5 ± 0.01	6.0 ± 0.1	6.2 ± 0.1	5.5 ± 0.3	6.3 ± 0.1	6.1 ± 0.1	–	19.3 ± 0.3
<i>B. anthracis</i>	ND	4.6 ± 0.1	5.6 ± 0.1	4.4 ± 0.3	2.1 ± 0.1	7.4 ± 0.2	–	16.7 ± 0.1
<i>E. coli</i>	ND	10.8 ± 0.1	8.1 ± 0.1	5.6 ± 0.3	6.8 ± 0.1	4.9 ± 0.1	–	18.5 ± 0.1
<i>K. pneumonia</i>	3.6 ± 0.01	10.3 ± 0.1	8.0 ± 0.2	7.2 ± 0.2	5.3 ± 0.1	6.2 ± 0.2	–	19.9 ± 0.2
<i>S. typhimurium</i>	ND	5.0 ± 0.1	13.3 ± 0.1	8.1 ± 0.2	5.0 ± 0.2	9.4 ± 0.1	–	19.6 ± 0.0
<i>P. aeuruginosa</i>	ND	4.1 ± 0.1	6.1 ± 0.2	5.3 ± 0.2	6.6 ± 0.2	8.8 ± 0.1	–	17.7 ± 0.0

Bold value represents the highest activity values observed

C. albicans (HG 392), *A. fumigatus* (HG 420), *A. niger* (ATCC 90028), *S. aureus*, (ATCC 25922), *S. faecalis* (ATCC 25925), *K. pneumonia* (ATCC 90028), *S. typhimurium* (ATCC 25927), *E. coli* (K 12), *P. aeuruginosa* (ATCC 25923), *B. anthracis*(QST 713)

Flu fluconazole, Am amoxicillin, ND not detected, – not done

* Values are mean \pm SD of three determinations

Acknowledgments

The authors wish to thank the National Commission for Science, Technology and Innovation (NACOSTI) for providing funds that enabled this research to be done. The Department of Chemistry, University of California at Berkeley and Technical University of Munich, Germany are thanked for NMR and ESI-MS analyses.

References

1. Abu Bakar NH, Sukari MA, Rahmani M, Sharif AM, Khalid K, Yusuf UK (2007) Chemical constituents from stem barks and roots of *Murraya koenigii* (Rutaceae). *Malays J Anal Sci* 11:173–176 [Google Scholar](#)
2. Bentjee H (1994) Kenyan trees, shrubs and lianas. National Museums of Kenya, Nairobi, pp 102–110 [Google Scholar](#)
3. Chakraborty DP, Bhattacharya PRS, Bhattacharya SP, Biswa AK (1978) Structure and synthesis of mukonine, a few carbazole alkaloids from *Murraya koenigii*. *Phytochemistry* 17:834–835 [CrossRef](#) [Google Scholar](#)
4. Chulay JD (1983) *Plasmodium falciparum*: assessment of in vitro growth by [3H]-hypoxanthine incorporation. *Exp Parasitol* 55:138–146 [CrossRef](#) [Google Scholar](#)
5. Desjardins RE, Canfield CJ, Haynes JD, Chulay JD (1979) Quantitative assessment of antimalarial activity *in vitro* by a semi-automated dilution technique. *Antimicrob Agents Chemother* 16:710–718
6. Dheeraf KG, Savita J, Pushpa D (2014) *Murraya koenigii* (L.) Spreng. An ethnobotanical, phytochemical and pharmacological review. *J Pharmacog Phytochem* 3:109–119 [Google Scholar](#)
7. Fiebig M, Pezzutp JM, Soejartp DD, Kinghorn AD (1985) Koeline, a further cytotoxic carbazole alkaloids from *Murraya koenigii*. *Phytochemistry* 24:3041–3043 [CrossRef](#) [Google Scholar](#)
8. Gbolade AA, Lukwa N, Kalenda D (2002) Guidelines for the studies on plant-based vector control agents. Obofemi Awolo University, Department of Pharmacognosy, Ile-Ile [Google Scholar](#)
9. Ito C, Itoigawa M, Nakao K, Murata T, Tsuboi M, Kaneda N, Furukawa H (2006) Induction of apoptosis by carbazole alkaloids isolated from *Murraya koenigii*. *Phytomedicine* 13:359–365 [CrossRef](#) [Google Scholar](#)
10. Kokwaro JO (2009) Medicinal plants of East Africa, 3rd edn. University of Nairobi Press, Nairobi, pp 134–137 [Google Scholar](#)
11. Kumar SS, Stephen J, Nthiya J (2014) Analysis of phytochemical status and antibacterial activities of *Alysicarpus bubleurifolius*-a valuable medicinal herb. *Am J Biol Life Sci* 2:146–149 [Google Scholar](#)
12. Lamers J, Buekert A, Makkar HPS, Von OM, Becker K (1996) Biomass production, feed and economic value of fodder weeds as by-products of millet cropping in a Sahelin farming system. *Exp Agric* 32:317–326 [CrossRef](#) [Google Scholar](#)
13. Mannetje L (2002) *Alysicarpus ovalifolius* (Schumach); J. Leonard record from protobase. In: Oyen LPA, Lemmens RHJ (eds) Plant resources of tropical Africa, vol 32. Bulletin du Jardin Botanique de l'État à Bruxelles, Bruxelles, pp 325–416 [Google Scholar](#)

14. McChesney J, Clark A, Silveira E (1991) Antibacterial diterpenes of *Croton sonderianus*. J Nat Prod 54:1625–1633 [CrossRef](#) [Google Scholar](#)
15. Mohammad S, Zahid H, Nirob KS, Manjural K, Nilufar N (2013) Antimicrobial activity of carbazole alkaloids from *Murraya koenigii* (L) Spreng leave. Int J Med Aromat Plants 3:131–135 [Google Scholar](#)
16. Nayak A, Mandal S, Banerji A, Banerji J (2010) Review on chemistry of *Murraya koenigii* Spreng (Rutaceae). J Chem Pharm Res 2:286–299 [Google Scholar](#)
17. Rameshkumar V, Umarajan KM (2013) Antibacterial activity of aqueous extract of *Alysicarpus longifolius* (Sperng), Wight & Arn against some pathogenic bacteria. J Modern Biotechnol 2:104–106 [Google Scholar](#)
18. Singh B, Sahu PM, Sharma MK (2002) Anti-inflammatory and anti-microbial activities of triterpenoids from *Strabilanthes callosus*. Phytomedicine 4:355–360 [CrossRef](#) [Google Scholar](#)
19. Trager W, Jensen JB (1976) Human malaria parasites in continuous culture. Science 193:673–675 [CrossRef](#) [Google Scholar](#)
20. Wojciechowski MF, Lavin M, Sanderson MJ (2004) A phylogeny of legumes (Leguminosae. Am J Botany 91(11):1846–1862 [CrossRef](#) [Google Scholar](#)
21. Yenesew A, Induli M, Derese S, Midiwo JO, Heydenreich M, Peter MG, Akala H, Wangui J, Liyala P, Waters NC (2004) Antiplasmodial flavonoids from the stem bark of *Erythrina abyssynica*. Phytochemistry 65:3029–3032 [CrossRef](#) [Google Scholar](#)

Copyright information

© The Korean Society for Applied Biological Chemistry 2015

Authors and Affiliations

- Angeline Atieno Ochung’
 - 1
- [Email author](#)
- Lawrence Arot Oyango Manguro
 - 1
- Phillip Okinda Owuor
 - 1
- Isaac Ogoche Jondiko
 - 1
- Regina Achieng’ Nyunja
 - 2
- Hosea Akala
 - 3
- Pauline Mwinzi
 - 3
- Sylvia Awino Opiyo

○ 1

1. 1.Department of ChemistryMaseno UniversityMasenoKenya
2. 2.Department of Biological SciencesJaramogi Oginga Odinga University of Science and TechnologyBondoKenya
3. 3.Kenya Medical Research Institute (KEMRI)KisumuKenya

About this article

Print ISSN

1738-2203

Online ISSN

2234-344X

Publisher Name

The Korean Society for Applied Biological Chemistry

[About this journal](#) [Reprints and Permissions](#)

Article actions

Export citation

- [.RIS Papers Reference Manager RefWorks Zotero](#)
- [.ENW EndNote](#)
- [.BIB BibTeX JabRef Mendeley](#)

Share article

- [Email](#)
- [Facebook](#)
- [Twitter](#)
- [LinkedIn](#)
- [Shareable link](#)

Article contents

- [Article](#)
- [Abstract](#)
- [Introduction](#)
- [Materials and methods](#)
- [Results and discussion](#)
- [Acknowledgments](#)
- [References](#)
- [Copyright information](#)
- [Authors and Affiliations](#)
- [About this article](#)

Cookies

We use cookies to improve your experience with our site. [More information](#)

Accept

Over 10 million scientific documents at your fingertips

Switch Edition

- [Academic Edition](#)
- [Corporate Edition](#)

- [Home](#)
- [Impressum](#)
- [Legal Information](#)
- [Contact Us](#)

© 2016 Springer International Publishing AG. Part of [Springer Nature](#).

Not logged in INASP - Kenya (2000510122) - Jaramogi Oginga Odinga University of Science & Technology (3000676541) 62.24.102.116