



JARAMOGI OGINGA ODINGA UNIVERSITY OF SCIENCE AND TECHNOLOGY

SCHOOL OF MATHEMATICS AND ACTUARIAL SCIENCE

UNIVERSITY EXAMINATION FOR DEGREE OF BACHELOR OF SCIENCE

ACTUARIAL

4TH YEAR SPECIAL RESITS – 2016

MAIN REGULAR

RESIT

COURSE CODE: SMA 202

COURSE TITLE: VECTOR ANALYSIS

EXAM VENUE: LAB 1

STREAM: (BSc. Actuarial)

DATE: 06/05/2016

EXAM SESSION: 2.00 – 4.00 PM

TIME: 2.00 HOURS

Instructions:

- 1. Answer question 1 (Compulsory) and ANY other 2 questions**
- 2. Candidates are advised not to write on the question paper.**
- 3. Candidates must hand in their answer booklets to the invigilator while in the examination room.**

QUESTION ONE (COMPULSORY)

- a) Given $\vec{r}_1 = 3\hat{i} - 2\hat{j} + \hat{k}$, $\vec{r}_2 = 2\hat{i} - 4\hat{j} - 3\hat{k}$, $\vec{r}_3 = -\hat{i} + 2\hat{j} + 2\hat{k}$, find the magnitude of $2\vec{r}_1 - 3\vec{r}_2 - 5\vec{r}_3$. (5 marks)
- b) Find a unit vector parallel to the resultant vectors $\vec{r}_1 = 2\hat{i} + 4\hat{j} - 5\hat{k}$, $\vec{r}_2 = \hat{i} + 2\hat{j} + 3\hat{k}$. (5 marks)
- c) Given the scalar field defined by $\phi(x, y, z) = 3x^2z - xy^3 + 5$, find ϕ at the point $(2, -2, 1)$. (4 marks)
- d) Find the angle between vectors $\vec{P} = 2\hat{i} + 2\hat{j} - \hat{k}$ and $\vec{Q} = 6\hat{i} - 3\hat{j} + 2\hat{k}$. (6 marks)
- e) If $\vec{S} = 2\hat{i} - 3\hat{j} - \hat{k}$ and $\vec{T} = \hat{i} + 4\hat{j} - 2\hat{k}$, find $|\vec{S} \times \vec{T}|$. (5 marks)
- f) Given $\vec{R} = \cos t\hat{i} + \sin t\hat{j} + 2t\hat{k}$, find $\left| \frac{d^2\vec{R}}{dt^2} \right|$. (5 marks)

QUESTION TWO

- a) Determine a unit vector parallel to the plane of $\vec{P} = 2\hat{i} - 6\hat{j} - 3\hat{k}$ and $\vec{Q} = 4\hat{i} + 3\hat{j} - \hat{k}$. (6 marks)
- b) If $\vec{A} = \hat{i} - 2\hat{j} - 3\hat{k}$, $\vec{B} = 2\hat{i} + \hat{j} - \hat{k}$ and $\vec{C} = \hat{i} + 3\hat{j} - 2\hat{k}$ find $|\vec{A} \times (\vec{B} \times \vec{C})|$. (6 marks)
- c) Evaluate $(2\hat{i} + \hat{j} - \hat{k}) \times (3\hat{i} - 2\hat{j} + 4\hat{k})$. (4 marks)
- d) Find the projection of the vector $2\hat{i} - 3\hat{j} + 6\hat{k}$ on the vector $\hat{i} + 2\hat{j} + 2\hat{k}$. (4 marks)

QUESTION THREE

- a) If $\vec{Q} = 5t^2\hat{i} + t\hat{j} - t^3\hat{k}$ and $R = \sin t\hat{i} - \cos t\hat{j}$, find $\frac{d}{dt}(\vec{Q} \times \vec{R})$. (6 marks)
- b) If $\vec{E} = (2x^2y - x^4)\hat{i} + (e^{xy} - y \sin x)\hat{j} + (x^2 \cos y)\hat{k}$, find $\frac{\partial^2 \vec{E}}{\partial y \partial x}$ at the point $(1, -1, 2)$. (6 marks)
- c) Find the unit tangent vector to any point on the curve $x = a \cos \omega t$, $y = a \sin \omega t$, $z = bt$ where a , b , ω are constants. (4 marks)
- d) If $\phi(x, y, z) = x^2yz$ and $\vec{F} = xz\hat{i} - xy^2\hat{j} + yz^2\hat{k}$, find $\frac{\partial^3}{\partial x \partial y \partial z}(\phi \vec{F})$ at the point $(2, -1, 1)$. (5 marks)

QUESTION FOUR

- a) Find $\nabla |r|^3$. (4 marks)
- b) If $\vec{F} = (3x^2y - z)\hat{i} + (xz^3 + y^4)\hat{j} - 2x^3z^2\hat{k}$, find $\nabla(\nabla \cdot \vec{F})$ at the point $(-1, 2, 0)$. (6 marks)

c) If $P = x^2yz$, $Q = xy - 3z^2$, find $\nabla \times [(\nabla P) \times (\nabla Q)]$. (6 marks)

d) Find the unit outward drawn normal to the surface $(x-1)^2 + y^2 + (z+1)^2 = 9$ at the point $(3, -1, 4)$. (4 marks)

QUESTION FIVE

a) If $\vec{P} = (4x^2 + 5y)\underline{i} - 12yz\underline{j} + 10xz^2\underline{k}$, evaluate $\int_c \vec{P} \cdot d\underline{r}$ from $(0, 0, 0)$ to $(1, 1, 1)$ along the

following path c :

i. $x = t, y = t^2, z = t^3$; (4 marks)

ii. the straight line from $(0, 0, 0)$ to $(1, 0, 0)$ then to $(1, 1, 0)$ and then to $(1, 1, 1)$; (4 marks)

iii. the straight line joining $(0, 0, 0)$ and $(1, 1, 1)$. (4 marks)

b) Evaluate $\iint_S \vec{F} \cdot \underline{n} ds$, where $\vec{F} = z\underline{i} + x\underline{j} - 3y^2z\underline{k}$ and S is the surface of the cylinder

$x^2 + y^2 = 16$ included in the first octant between $z = 0$ and $z = 5$. (8 marks)