

QUESTION 1 (30 MARKS)

a)

- i. State **TWO** postulates of Quantum mechanics(**1 mark**)
- ii. Show that the solution of the time-dependent Schrödinger equation takes the form $\Psi(\vec{r}, t) = \psi(\vec{r})u(0)e^{\frac{-iEt}{\hbar}}$ where the symbols have their usual meanings.(**3 marks**)

b) Using Dirac's notation of eigenfunctions, state the Riesz variational principle. **(1 mark)**

c) Define the following terms as used in Quantum mechanics.

- i. Spin-orbit coupling(**1 mark**)
- ii. Schrödinger picture(**1 mark**)
- iii. Heisenberg picture(**1 mark**)
- iv. Interaction picture **(1 mark)**

d) Show that the x-component of the orbital angular momentum is given by

$$\hat{L}_x = i\hbar \left(\sin \phi \frac{\partial}{\partial \theta} + \cos \phi \frac{\partial}{\partial \phi} \right) \text{ (4 marks)}$$

e) Show that the time-independent Schrödinger equation of a hydrogenic atom

is given by $\left[-\frac{\hbar^2}{2\mu} \nabla^2 - \frac{Ze^2}{4\pi\epsilon_0 r} \right] \psi(\vec{r}) = E\psi(\vec{r})$ where the symbols have their usual

meanings. **(4 marks)**

f) Distinguish between time-independent perturbation theory and time-dependent perturbation theory. **(2 marks)**

g) State the selection rules for allowed transitions in hydrogen atom. **(2 marks)**

h) Account for the Pauli exclusion principle for fermions.**(2 marks)**

i) Derive the Heisenberg's equation of motion. **(4 marks)**

j) The spin-up and spin-down state vectors of an electron are respectively

defined by $|u\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}; |d\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Write down the Hermitian conjugate state

vectors $\langle u|; \langle d|$ and show that they satisfy the orthonormality relations

$$\langle u|u\rangle = \langle d|d\rangle = 1; \langle u|d\rangle = \langle d|u\rangle = 0 \text{ (3 marks)}$$

SECTION B

Attempt any TWO questions in this section.

QUESTION 2 (20 MARKS)

(i) Express the Hamiltonian of a one-dimensional linear harmonic oscillator in

the form $\hat{H} = \hbar\omega\left(a^+a + \frac{1}{2}\right)$ where \hat{a} , \hat{a}^+ are the usual annihilation and creation

operators which must be defined in the derivation. **(7 marks)**

(ii) Calculate the energy spectrum of the oscillator in the number state $|n\rangle$.

(4 marks)

(iii) Show that the ground state of the oscillator is a minimum uncertainty state,

hence give the physical interpretation of such a state. **(9 marks)**

QUESTION 3 (20 MARKS)

a) (i) Matrix operators for the angular momentum operators can be defined by

$$\hat{L}_x = \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}; \hat{L}_y = \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}; \hat{L}_z = \hbar \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \text{ Determine the}$$

commutator $[\hat{L}_x, \hat{L}_y]$. **(3 marks)**

(ii) Show that the ladder operator \hat{L}_+ , of the angular momentum in spherical

coordinates takes the form $\hat{L}_+ = \hbar e^{i\phi} \left(\frac{\partial}{\partial \theta} + i \cot \theta \frac{\partial}{\partial \phi} \right)$ **(7 marks)**

b) A particle moves in the one-dimensional potential defined by

$$V(x) = \begin{cases} V_0 \cos\left(\frac{\pi x}{2a}\right) & |x| \leq a \\ \infty & |x| > a \end{cases}. \text{ By treating the potential as a perturbation, obtain}$$

the first order energy correction, given that the unperturbed eigen function is

$$u_n = \begin{cases} \frac{1}{\sqrt{a}} \cos\left(\frac{n\pi x}{2a}\right) & n = \text{odd} \\ \frac{1}{\sqrt{a}} \sin\left(\frac{n\pi x}{2a}\right) & n = \text{even} \end{cases}. \quad (10 \text{ marks})$$

QUESTION 4 (20 MARKS)

a) Using $u(r) = rR(r)$, the radial equation for a one-electron atom is obtained in the form $\left(-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} + \frac{\hbar^2 l(l+1)}{2\mu r^2} - \frac{Ze^2}{4\pi\epsilon_0 r}\right)u(r) = Eu(r)$ where the symbols have their usual meanings.

(i) By completing the square of the effective potential, determine the

quantized orbit energy in the form $E_{l+1} = \frac{-\mu Z^2 e^4}{2(4\pi\epsilon_0)^2 \hbar^2 (l+1)^2}$. (5 marks)

(ii) Show that the radial equation can be factorized in the form

$$\left(\frac{d}{dr} + K_{l+1}(r)\right)\left(-\frac{d}{dr} + K_{l+1}(r)\right)u(r) = \frac{2\mu}{\hbar^2}(E - E_{l+1})u(r)$$
 where the parameter $K_{l+1}(r)$

must be defined in the derivation. (8 marks)

b) Determine the highest quantized orbit solution of the factorized radial equation in a(ii) above in the form $u_n(r) = Ar^n e^{\frac{-Zr}{a_0 n}}$: $n = 1, 2, \dots$ where a_0 is the Bohr radius. (7 marks)

QUESTION 5 (20 MARKS)

a) A two-level system described by the wave function

$$\psi(t) = c_a(t)\psi_a e^{-iE_a \frac{t}{\hbar}} + c_b(t)\psi_b e^{-iE_b \frac{t}{\hbar}}$$
 experiences a time-dependent perturbation.

Suppose the system is in state ψ_a initially, derive the expressions for the

first order approximations of probability amplitudes, $c_a^1(t)$ and $c_b^1(t)$.
(12 marks)

b) If the perturbation in 5 (a) above is of the form $H'_{ab} = V_{ab} \cos \omega t$, show that the transition probability is given by $P_{a \rightarrow b} = \frac{|V_{ab}|^2}{\hbar^2} \frac{\sin^2(\omega_0 - \omega)t}{(\omega_0 - \omega)^2}$ **(8 marks)**