



JARAMOGI OGINGA ODINGA UNIVERSITY OF SCIENCE AND TECHNOLOGY
SCHOOL OF BIOLOGICAL, PHYSICAL, MATHEMATICS AND ACTUARIAL
SCIENCE
UNIVERSITY EXAMINATION FOR DEGREE OF BACHELOR OF SCIENCE
ACTUARIAL
3RD YEAR 1ST SEMESTER 2022/2023 ACADEMIC YEAR
MAIN REGULAR

COURSE CODE:WAB 2309

COURSE TITLE:THEORY OF ESTIMATION

EXAM VENUE: LAB 17

STREAM: (BSc. Actuarial)

DATE: 15/12/2022

EXAM SESSION: 15.00-17.00PM

TIME: 2.00 HOURS

Instructions:

- 1. Answer question 1 (Compulsory) and ANY other 2 questions**
- 2. Candidates are advised not to write on the question paper.**
- 3. Candidates must hand in their answer booklets to the invigilator while in the examination room.**

QUESTION ONE (20 MARKS)

- a) Explain clearly the following terms as used in theory of estimation.
- Sufficiency
 - Weak Consistency
 - Completeness
 - Uniformly Minimum Variance Unbiased Estimator.
 - Most Efficient Estimator
 - Unbiasedness
- [6 Marks]
- b) Let $X_1, X_2, X_3, \dots, X_n$ be a random sample of size n from the Poisson population with parameter λ . Show that $T_n = \frac{\bar{x}}{1 + \frac{1}{n}}$ is consistent for λ . [6 Marks]
- c) Let X_1, X_2, \dots, X_n be a random sample of size n from the exponential distribution with pdf $f(x; \theta) = \frac{1}{\theta} e^{-x/\theta}$. Show that \bar{X} is unbiased for θ . [6 Marks]
- d) Given two estimators of population mean (μ) as $T_1 = \frac{(X_1 + 2X_2 + X_3)}{4}$ and $T_2 = \frac{(X_1 + X_2 + X_3)}{3}$ where X_1, X_2 and X_3 are from $N(\mu, \sigma^2)$ distribution. Prove that T_2 is more efficient than T_1 . [6 Marks]
- e) Let X_1, X_2, \dots, X_n be iid *Uniform* $[a, b]$ random variables with a known. Find an unbiased estimator for b . [6 Marks]

QUESTION TWO (20 MARKS)

- a) Let X_1, X_2, \dots, X_n be iid random variables from the $N(\mu, \sigma^2)$ distribution. Find the maximum likelihood estimators of μ and σ^2 [11 Marks]
- b) Let X_1, X_2, \dots, X_n be iid random variables from a uniform distribution on the interval $(\theta - 1, \theta + 1)$.
- Find the method of moments estimator of θ .
 - Is the obtained estimator unbiased for θ ?
- [9 Marks]

QUESTION THREE (20 MARKS)

- a) Use the Lehmann Scheffe method of construction of minimal sufficient statistics to find the minimal sufficient statistic for $\theta = (\mu, \sigma^2)$ given X_1, X_2, \dots, X_n are iid random variables from (μ, σ^2) . [9 Marks]
- b) Let X_1, X_2, \dots, X_n be a random sample of size n from $N(\mu, \sigma^2)$. Find Fisher's information $I(\mu)$ necessary for estimation of the Population mean (μ) hence the associated Cramer-Rao lower bound. [11.Marks]

QUESTION FOUR (20 MARKS)

Let X_1, X_2, \dots, X_n be a random sample from a population having probability density function;

$$f(x) = \begin{cases} \frac{2}{\theta^2}(\theta - x), & 0 < x < \theta \\ 0, & \text{Otherwise} \end{cases}$$

Let $T = \frac{3}{2}\bar{x}$ be an estimator of θ . Find the Root Mean Squared Error of the estimator T

[20 Marks]

QUESTION FIVE (20 MARKS)

- a) Let X_1, X_2, \dots, X_n be iid Poisson random variables with $f(x, \theta) = \frac{e^{-\theta}\theta^x}{x!} x = 0, 1, 2, \dots$
 Show that the maximum likelihood estimator for θ is \bar{X} . **[8 Marks]**

- b) Let X_1, X_2, \dots, X_n be a random sample of n observations from a population having p.d.f

$$f(x) = \begin{cases} \frac{2x}{\theta^2}, & 0 \leq x \leq \theta \\ 0, & \text{otherwise} \end{cases}$$

Check if $T = \frac{3\bar{x}}{2}$ is consistent for θ

[12 Marks]