



**JARAMOGI OGINGA ODINGA UNIVERSITY OF SCIENCE AND TECHNOLOGY
SCHOOL OF BIOLOGICAL AND PHYSICAL SCIENCE
UNIVERSITY EXAMINATION FOR DEGREE OF BACHELOR OF EDUCATION
(ARTS/SCIENCE/SNE)
4TH YEAR 2ND SEMESTER 2013/2014 ACADEMIC YEAR
MAIN SCHOOL BASED**

COURSE CODE: SCH 411

COURSE TITLE: ORGANIC STEREO CHEMISTRY

EXAM VENUE: LAB 2

STREAM: (BEd Arts/science)

DATE: 29/08/14

EXAM SESSION: 9.00 – 11.00 A M

TIME: 2.00 HOURS




Instructions:

- 1. Answer ALL questions in section A and ANY other 2 questions in section B**
- 2. Candidates are advised not to write on the question paper.**
- 3. Candidates must hand in their answer booklets to the invigilator while in the examination room.**

SECTION A:

QUESTION 1

- a) Explain the following terms; (10 marks)
- Stereochemistry
 - Optical activity
 - Racemic mixture
 - Molecular geometry
 - Chirality
- b) Discuss the principles underlying the naming of enantiomers. (5 marks)
- c) Account for the following observations; (6 marks)
- 1,3-dimethylcyclohexane exists only in three rather than four isomers though it has two stereogenic centres.
 - A pair of enantiomers has identical infra red spectra, indexes of reflection, solubilities and reaction kinetics in ordinary solvents.
 - Molecular geometries can be specified in terms of bond lengths, bond angles and torsional angles.
- d) The following are stereorepresentations for the three stereoisomers of 2,3-butanediol. The carbons are numbered beginning from left as shown in (1).




- Assign an *R* or *S* configuration to each chiral centre. (3 marks)
 - Which are enantiomers? (1 mark)
 - Which is the meso compound? (1 mark)
 - Which are diastereomers? (1 mark)
- e) Draw the stereoisomers that exist for 1,2-cyclopentanediol molecule. (3 marks)

SECTION B (40 MARKS):

QUESTION 2


- a) Define each of the following terms: (10 marks)
- Diastereomers
 - Stereogenic centre
 - Meso compound
 - Enantiometrically pure substances
 - Solvolysis reaction

- b) Consider the reaction below: (6 marks)



How many stereoisomers of the product are possible? Draw them. Are the products optically active?

- c) What products are formed from the following reaction? Are the product(s) optically active? (4 marks)



QUESTION 3

- a) Distinguish the following terms; (4 marks)

- Stereoisomers and constitutional isomers
- Stereospecific and regiospecific reactions

- b) At 24°C, a sample of S-2-iodobutane whose specific rotation is: $[\alpha]_D^{24} = -22.4^\circ$ was put in a 1 dm vial of solution of 1 gml⁻¹ showed an optical rotation of +3.975°.

- What is the optical purity? (4 marks)
- What is the enantiomeric excess? (4 marks)

- c) Which of the following compounds are chiral? (4 marks)

- 2,4-dimethylheptane
- 5-ethyl-3,3-dimethyloctane
- Cis-1,4-dichlorocyclohexane
- 4,5-dimethyl-2,6-octadiyne

- d) What is the relationship between the specific rotations of; (4 marks)


- (2R,3R)-dichloropentane and (2S,3S)-dichloropentane
- (2R,3S)-dichloropentane and (2S,3R)-dichloropentane.

QUESTION 4

- a) Draw the tetrahedral representations of the following compounds; (4 marks)

- (S)-2-chlorobutane
- (R)-chloro-1-pentene

- b) Assign R or S configurations to the chiral centers of ascorbic acid. (2 marks)



- b) Give an example of a compound with the following molecular structure. (8 marks)
- Linear
 - Trigonal planar
 - Bent
 - Tetrahedral
- c) The specific rotation of progesterone a female hormone is $+172^\circ$. Calculate the observed rotation for a solution prepared by dissolving 300 mg of progesterone in 15.0 ml of dioxane and placing it in a sample tube of 10.0 cm long. (6 marks)

QUESTION 5

- a) Giving relevant examples, describe how racemic mixtures can be resolved using any **TWO** of the following methods.
- Use of chiral bases or acids (6 marks)
 - Chromatography (6 marks)
 - Use of enzymes (6 marks)
- b) Many drugs are isolated directly from plants and bacteria while others are obtained by modification of naturally occurring compounds. Such compounds are usually chiral and are found as a single enantiomer rather than a racemic mixture. Giving relevant names and structures, discuss application of four such drugs. (8 marks)